arXiv:1301.4476v1 [math.CA] 17 Jan 2013

TWO TRANSFORMATION FORMULAS FOR ${}_7\psi_7$-SERIES

[1]CHUANAN WEI, [2]QINGLUN YAN, [3]DIANXUAN GONG*

[1]*Department of Information Technology*
Hainan Medical College, Haikou 571199, China
[2]*College of Mathematics and Physics*
Nanjing University of Posts and Telecommunications, Nanjing 210046, China
[3]*College of Sciences*
Hebei United University, Tangshan 063009, China

Abstract. In terms of Cauchy's method, we establish two transformation formulas for ${}_7\psi_7$-series. Surprisingly, they include the direct nonterminating generalizations of four summation formulas for bilateral q-series as special cases.

1. Introduction

For two complex numbers q and x, define the q-shifted factorial by

$$(x;q)_n=\begin{cases}\prod_{i=0}^{n-1}(1-xq^i), & \text{when} \quad n>0;\\ 1, & \text{when} \quad n=0;\\ \frac{1}{\prod_{j=n}^{-1}(1-xq^j)}, & \text{when} \quad n<0.\end{cases}$$

When $|q|<1$, the q-shifted factorial of infinite order reads as

$$(x;q)_\infty=\prod_{k=0}^{\infty}(1-xq^k).$$

For simplifying the expressions, we shall use the following compact notations:

$$\left[\begin{matrix}a,b,\cdots,c\\ \alpha,\beta,\cdots,\gamma\end{matrix}\Big|\, q\right]_n=\frac{(a;q)_n(b;q)_n\cdots(c;q)_n}{(\alpha;q)_n(\beta;q)_n\cdots(\gamma;q)_n},$$

$$\left[\begin{matrix}a,b,\cdots,c\\ \alpha,\beta,\cdots,\gamma\end{matrix}\Big|\, q\right]_\infty=\frac{(a;q)_\infty(b;q)_\infty\cdots(c;q)_\infty}{(\alpha;q)_\infty(\beta;q)_\infty\cdots(\gamma;q)_\infty}.$$

Following Gasper and Rahman [8], define the unilateral q-series by

$${}_{1+r}\phi_s\left[\begin{matrix}a_0, & a_1, & \cdots,a_r\\ & b_1, & \cdots,b_s\end{matrix}\Big|\, q;z\right]=\sum_{k=0}^{\infty}\left[\begin{matrix}a_0,a_1,\cdots,a_r\\ q,\,b_1,\,\cdots,b_s\end{matrix}\Big|\, q\right]_k\left\{(-1)^kq^{\binom{k}{2}}\right\}^{s-r}z^k.$$

Then Bailey's four-term transformation formula (cf. [8, p. 57]) can be written as

---

2010 Mathematics Subject Classification: Primary 05A19 and Secondary 33D15.

Key words and phrases. Cauchy's method; Unilateral q-series; Bilateral q-series.

*Corresponding author. *Email addresses*: weichuanan@yahoo.com.cn (C. Wei), yanqinglun@yahoo.com.cn (Q. Yan), gongdianxuan@yahoo.com.cn (D. Gong).

$$\begin{aligned}
&{}_{10}\phi_9\left[\begin{matrix}a, q\sqrt{a}, -q\sqrt{a}, b, c, d, e, f, g, h\\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq/d, aq/e, aq/f, aq/g, aq/h\end{matrix}\middle| q;q\right]\\
&+\left[\begin{matrix}aq,b/a,c,d,e,f,g,h,bq/c,bq/d,bq/e,bq/f,bq/g,bq/h\\ b^2q/a,a/b,aq/c,aq/d,aq/e,aq/f,aq/g,aq/h,bc/a,bd/a,be/a,bf/a,bg/a,bh/a\end{matrix}\middle| q\right]_\infty\\
&\times {}_{10}\phi_9\left[\begin{matrix}b^2/a, qb/\sqrt{a}, -qb/\sqrt{a}, b, bc/a, bd/a, be/a, bf/a, bg/a, bh/a\\ b/\sqrt{a}, -b/\sqrt{a}, bq/a, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h\end{matrix}\middle| q;q\right]\\
&=\left[\begin{matrix}aq, b/a, \lambda q/f, \lambda q/g, \lambda q/h, bf/\lambda, bg/\lambda, bh/\lambda\\ \lambda q, b/\lambda, aq/f, aq/g, aq/h, bf/a, bg/a, bh/a\end{matrix}\middle| q\right]_\infty\\
&\times {}_{10}\phi_9\left[\begin{matrix}\lambda, q\sqrt{\lambda}, -q\sqrt{\lambda}, b, \lambda c/a, \lambda d/a, \lambda e/a, f, g, h\\ \sqrt{\lambda}, -\sqrt{\lambda}, \lambda q/b, aq/c, aq/d, aq/e, \lambda q/f, \lambda q/g, \lambda q/h\end{matrix}\middle| q;q\right]\\
&+\left[\begin{matrix}aq,b/a,f,g,h,bq/f,bq/g,bq/h,\lambda c/a,\lambda d/a,\lambda e/a,abq/\lambda c,abq/\lambda d,abq/\lambda e\\ b^2q/\lambda,\lambda/b,aq/c,aq/d,aq/e,aq/f,aq/g,aq/h,bc/a,bd/a,be/a,bf/a,bg/a,bh/a\end{matrix}\middle| q\right]_\infty\\
&\times {}_{10}\phi_9\left[\begin{matrix}b^2/\lambda, qb/\sqrt{\lambda}, -qb/\sqrt{\lambda}, b, bc/a, bd/a, be/a, bf/\lambda, bg/\lambda, bh/\lambda\\ b/\sqrt{\lambda}, -b/\sqrt{\lambda}, bq/\lambda, abq/\lambda c, abq/\lambda d, abq/\lambda e, bq/f, bq/g, bq/h\end{matrix}\middle| q;q\right] \quad (1)
\end{aligned}$$

where $\lambda = qa^2/cde$ and $q^2a^3 = bcdefgh$.

Following the same book, define the bilateral q-series by

$$ {}_r\psi_s\left[\begin{matrix}a_1, & \cdots, a_r\\ b_1, & \cdots, b_s\end{matrix}\middle| q;z\right] = \sum_{k=-\infty}^{\infty}\left[\begin{matrix}a_1,\cdots, a_r\\ b_1,\cdots, b_s\end{matrix}\middle| q\right]_k \left\{(-1)^k q^{\binom{k}{2}}\right\}^{s-r} z^k. $$

Then two nice ${}_3\psi_3$-series identities due to Bailey [1, Equations (2.2) and (2.3)] can be stated as follows:

$$ {}_3\psi_3\left[\begin{matrix}b, c, d\\ q/b, q/c, q/d\end{matrix}\middle| q;\frac{q}{bcd}\right] = \left[\begin{matrix}q, q/bc, q/bd, q/cd\\ q/b, q/c, q/d, q/bcd\end{matrix}\middle| q\right]_\infty, \quad (2)$$

$$ {}_3\psi_3\left[\begin{matrix}b, c, d\\ q^2/b, q^2/c, q^2/d\end{matrix}\middle| q;\frac{q^2}{bcd}\right] = \left[\begin{matrix}q, q^2/bc, q^2/bd, q^2/cd\\ q^2/b, q^2/c, q^2/d, q^2/bcd\end{matrix}\middle| q\right]_\infty, \quad (3)$$

where the convergent conditions for (2) and (3) are $|q/bcd| < 1$ and $|q^2/bcd| < 1$ respectively. The finite forms of them read as

$$\begin{aligned}
&{}_5\psi_5\left[\begin{matrix}b, c, d, q^{n+1}/bcd, q^{-n}\\ q/b, q/c, q/d, bcd/q^n, q^{n+1}\end{matrix}\middle| q;q\right]\\
&=\left[\begin{matrix}q, q/bc, q/bd, q/cd\\ q/b, q/c, q/d, q/bcd\end{matrix}\middle| q\right]_n, \quad (4)\\
&{}_5\psi_5\left[\begin{matrix}b, c, d, q^{n+3}/bcd, q^{-n}\\ q^2/b, q^2/c, q^2/d, bcd/q^{n+1}, q^{n+2}\end{matrix}\middle| q;q\right]\\
&=\left[\begin{matrix}q^2, q^2/bc, q^2/bd, q^2/cd\\ q^2/b, q^2/c, q^2/d, q^2/bcd\end{matrix}\middle| q\right]_n (1-q), \quad (5)
\end{aligned}$$

where (4) and (5) are due to Jackson [4, Equation (1)] and Bailey [1, Equation (3.2)] respectively.

Recently, Cauchy's method has been utilized to study the bilateral q-series in [2], [5], [6], [7], [10] and [11]. Inspired by these work, we shall derive two transformation formulas for ${}_7\psi_7$-series, which include the direct nonterminating generalizations of (2), (3), (4) and (5) as special cases, from (1) in the same way.

2. Two transformation formulas for $_7\psi_7$-series

Theorem 1. *For $u = q/cde$ and $q^2 = bcdefgh$, there holds the transformation formula for $_7\psi_7$-series:*

$$
\begin{aligned}
&{}_7\psi_7\left[\begin{matrix} b, c, d, e, f, g, h \\ q/b, q/c, q/d, q/e, q/f, q/g, q/h \end{matrix} \middle| q; q\right] \\
&+ \left[\begin{matrix} q, b, c, d, e, f, g, h, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \\ b^2q, 1/b, q/c, q/d, q/e, q/f, q/g, q/h, bc, bd, be, bf, bg, bh \end{matrix} \middle| q\right]_\infty \\
&\times {}_8\phi_7\left[\begin{matrix} b^2, -qb, bc, bd, be, bf, bg, bh \\ -b, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \end{matrix} \middle| q; q\right] \\
&= \left[\begin{matrix} q, b, uq/f, uq/g, uq/h, bf/u, bg/u, bh/u \\ uq, b/u, q/f, q/g, q/h, bf, bg, bh \end{matrix} \middle| q\right]_\infty \\
&\times {}_{10}\phi_9\left[\begin{matrix} u, q\sqrt{u}, -q\sqrt{u}, b, uc, ud, ue, f, g, h \\ \sqrt{u}, -\sqrt{u}, uq/b, q/c, q/d, q/e, uq/f, uq/g, uq/h \end{matrix} \middle| q; q\right] \\
&+ \left[\begin{matrix} q, b, f, g, h, bq/f, bq/g, bq/h, uc, ud, ue, bq/uc, bq/ud, bq/ue \\ b^2q/u, u/b, q/c, q/d, q/e, q/f, q/g, q/h, bc, bd, be, bf, bg, bh \end{matrix} \middle| q\right]_\infty \\
&\times {}_{10}\phi_9\left[\begin{matrix} b^2/u, qb/\sqrt{u}, -qb/\sqrt{u}, b, bc, bd, be, bf/u, bg/u, bh/u \\ b/\sqrt{u}, -b/\sqrt{u}, bq/u, bq/uc, bq/ud, bq/ue, bq/f, bq/g, bq/h \end{matrix} \middle| q; q\right].
\end{aligned}
$$

Proof. The limiting case $a \to 1$ of (1) leads to the following equation:

$$
\begin{aligned}
&\left\{1 + \sum_{k=1}^{\infty}\{1+q^k\}\left[\begin{matrix} b, c, d, e, f, g, h \\ q/b, q/c, q/d, q/e, q/f, q/g, q/h \end{matrix} \middle| q\right]_k q^k\right\} \\
&+ \left[\begin{matrix} q, b, c, d, e, f, g, h, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \\ b^2q, 1/b, q/c, q/d, q/e, q/f, q/g, q/h, bc, bd, be, bf, bg, bh \end{matrix} \middle| q\right]_\infty \\
&\times {}_8\phi_7\left[\begin{matrix} b^2, -qb, bc, bd, be, bf, bg, bh \\ -b, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \end{matrix} \middle| q; q\right] \\
&= \left[\begin{matrix} q, b, uq/f, uq/g, uq/h, bf/u, bg/u, bh/u \\ uq, b/u, q/f, q/g, q/h, bf, bg, bh \end{matrix} \middle| q\right]_\infty \\
&\times {}_{10}\phi_9\left[\begin{matrix} u, q\sqrt{u}, -q\sqrt{u}, b, uc, ud, ue, f, g, h \\ \sqrt{u}, -\sqrt{u}, uq/b, q/c, q/d, q/e, uq/f, uq/g, uq/h \end{matrix} \middle| q; q\right] \\
&+ \left[\begin{matrix} q, b, f, g, h, bq/f, bq/g, bq/h, uc, ud, ue, bq/uc, bq/ud, bq/ue \\ b^2q/u, u/b, q/c, q/d, q/e, q/f, q/g, q/h, bc, bd, be, bf, bg, bh \end{matrix} \middle| q\right]_\infty \\
&\times {}_{10}\phi_9\left[\begin{matrix} b^2/u, qb/\sqrt{u}, -qb/\sqrt{u}, b, bc, bd, be, bf/u, bg/u, bh/u \\ b/\sqrt{u}, -b/\sqrt{u}, bq/u, bq/uc, bq/ud, bq/ue, bq/f, bq/g, bq/h \end{matrix} \middle| q; q\right],
\end{aligned}
$$

where $u = q/cde$ and $q^2 = bcdefgh$. The first term on the left hand side can be reformulated as

$$
\begin{aligned}
&1 + \sum_{k=1}^{\infty}\{1+q^k\}\left[\begin{matrix} b, c, d, e, f, g, h \\ q/b, q/c, q/d, q/e, q/f, q/g, q/h \end{matrix} \middle| q\right]_k q^k \\
&= 1 + \sum_{k=1}^{\infty}\left[\begin{matrix} b, c, d, e, f, g, h \\ q/b, q/c, q/d, q/e, q/f, q/g, q/h \end{matrix} \middle| q\right]_k q^k \\
&\quad + \sum_{k=1}^{\infty}\left[\begin{matrix} b, c, d, e, f, g, h \\ q/b, q/c, q/d, q/e, q/f, q/g, q/h \end{matrix} \middle| q\right]_k q^{2k}
\end{aligned}
$$

$$\begin{aligned}
&= 1+\sum_{k=1}^{\infty}\left[\begin{matrix} b,c,d,e,f,g,h \\ q/b,q/c,q/d,q/e,q/f,q/g,q/h \end{matrix}\Big| q\right]_k q^k \\
&\quad+\sum_{k=-\infty}^{-1}\left[\begin{matrix} b,c,d,e,f,g,h \\ q/b,q/c,q/d,q/e,q/f,q/g,q/h \end{matrix}\Big| q\right]_{-k} q^{-2k} \\
&= 1+\sum_{k=1}^{\infty}\left[\begin{matrix} b,c,d,e,f,g,h \\ q/b,q/c,q/d,q/e,q/f,q/g,q/h \end{matrix}\Big| q\right]_k q^k \\
&\quad+\sum_{k=-\infty}^{-1}\left[\begin{matrix} b,c,d,e,f,g,h \\ q/b,q/c,q/d,q/e,q/f,q/g,q/h \end{matrix}\Big| q\right]_{k} q^{k} \\
&= {}_7\psi_7\left[\begin{matrix} b,c,d,e,f,g,h \\ q/b,q/c,q/d,q/e,q/f,q/g,q/h \end{matrix}\Big| q;q\right].
\end{aligned}$$

Therefore, we obtain Theorem 1 to complete the proof. ☐

Taking $b=q^{n+2}/cdefg, h=q^{-n}$ in Theorem 1 and then letting $n\to\infty$, we get the following result under the replacements $e\to b$ and $g\to e$.

Corollary 2. *For $\mu=q/bcd$ and $\max\{|q^2/bcdef|,|q/ef|\}<1$, there holds the transformation formula between a ${}_5\psi_5$-series and a ${}_8\phi_7$-series:*

$$\begin{aligned}
&{}_5\psi_5\left[\begin{matrix} b,c,d,e,f \\ q/b,q/c,q/d,q/e,q/f \end{matrix}\Big| q;\frac{q^2}{bcdef}\right]=\left[\begin{matrix} q,q/ef,\mu q/e,\mu q/f \\ q/e,q/f,\mu q/ef,\mu q \end{matrix}\Big| q\right]_\infty \\
&\times\ {}_8\phi_7\left[\begin{matrix} \mu,q\sqrt{\mu},-q\sqrt{\mu},\mu b,\mu c,\mu d,e,f \\ \sqrt{\mu},-\sqrt{\mu},q/b,q/c,q/d,\mu q/e,\mu q/f \end{matrix}\Big| q;\frac{q}{ef}\right].
\end{aligned}$$

When $c=q/d$, Corollary 2 reduces to (2) under the substitutions $e\to c$ and $f\to d$.

Setting $c=q/d$ in Theorem 1, we attain the following result under the replacements $g\to c$ and $h\to d$.

Corollary 3. *For $q=bcdef$, there holds the transformation formula between a ${}_5\psi_5$-series and a ${}_6\phi_5$-series:*

$$\begin{aligned}
&{}_5\psi_5\left[\begin{matrix} b,c,d,e,f \\ q/b,q/c,q/d,q/e,q/f \end{matrix}\Big| q;q\right]={}_6\phi_5\left[\begin{matrix} b^2,-qb,bc,bd,be,bf \\ -b,bq/c,bq/d,bq/e,bq/f \end{matrix}\Big| q;q\right] \\
&\times\ b\left[\begin{matrix} q,bq,c,d,e,f,bq/c,bq/d,bq/e,bq/f \\ b^2q,q/b,q/c,q/d,q/e,q/f,bc,bd,be,bf \end{matrix}\Big| q\right]_\infty \\
&+\ \left[\begin{matrix} q,b,q/cd,q/ce,q/cf,q/de,q/df,q/ef \\ q/c,q/d,q/e,q/f,bc,bd,be,bf \end{matrix}\Big| q\right]_\infty.
\end{aligned}$$

When $e=q^{n+1}/bcd$ and $f=q^{-n}$, Corollary 3 reduces to (4) exactly.

Theorem 4. *For $v=q^3/cde$ and $q^5=bcdefgh$, there holds the transformation formula for ${}_7\psi_7$-series:*

$$\begin{aligned}
&{}_7\psi_7\left[\begin{matrix} b,c,d,e,f,g,h \\ q^2/b,q^2/c,q^2/d,q^2/e,q^2/f,q^2/g,q^2/h \end{matrix}\Big| q;q\right] \\
&+\ \left[\begin{matrix} q,b/q,c,d,e,f,g,h,bq/c,bq/d,bq/e,bq/f,bq/g,bq/h \\ b^2,q/b,q^2/c,q^2/d,q^2/e,q^2/f,q^2/g,q^2/h,bc/q,bd/q,be/q,bf/q,bg/q,bh/q \end{matrix}\Big| q\right]_\infty
\end{aligned}$$

$$
\begin{aligned}
&\times {}_9\phi_8\left[\begin{matrix} b^2/q, b\sqrt{q}, -b\sqrt{q}, bc/q, bd/q, be/q, bf/q, bg/q, bh/q \\ b/\sqrt{q}, -b/\sqrt{q}, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \end{matrix}\middle|\, q;q\right]\\
=\;& \left[\begin{matrix} q, b/q, vq/f, vq/g, vq/h, bf/v, bg/v, bh/v \\ vq, b/v, q^2/f, q^2/g, q^2/h, bf/q, bg/q, bh/q \end{matrix}\middle|\, q\right]_\infty\\
&\times {}_{10}\phi_9\left[\begin{matrix} v, q\sqrt{v}, -q\sqrt{v}, b, vc/q, vd/q, ve/q, f, g, h \\ \sqrt{v}, -\sqrt{v}, vq/b, q^2/c, q^2/d, q^2/e, vq/f, vq/g, vq/h \end{matrix}\middle|\, q;q\right]\\
+\;& \left[\begin{matrix} q,b/q,f,g,h,bq/f,bq/g,bq/h,vc/q,vd/q,ve/q,bq^2/vc,bq^2/vd,bq^2/ve \\ b^2q/v,v/b,q^2/c,q^2/d,q^2/e,q^2/f,q^2/g,q^2/h,bc/q,bd/q,be/q,bf/q,bg/q,bh/q \end{matrix}\middle|\, q\right]_\infty\\
&\times {}_{10}\phi_9\left[\begin{matrix} b^2/v, qb/\sqrt{v}, -qb/\sqrt{v}, b, bc/q, bd/q, be/q, bf/v, bg/v, bh/v \\ b/\sqrt{v}, -b/\sqrt{v}, bq/v, bq^2/vc, bq^2/vd, bq^2/ve, bq/f, bq/g, bq/h \end{matrix}\middle|\, q;q\right].
\end{aligned}
$$

Proof. The limiting case $a \to q$ of (1) leads to the following equation:

$$
\begin{aligned}
&\sum_{k=0}^{\infty}\{1-q^{1+2k}\}\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\middle|\, q\right]_k q^k\\
+\;& \left[\begin{matrix} q,b/q,c,d,e,f,g,h,bq/c,bq/d,bq/e,bq/f,bq/g,bq/h \\ b^2,q/b,q^2/c,q^2/d,q^2/e,q^2/f,q^2/g,q^2/h,bc/q,bd/q,be/q,bf/q,bg/q,bh/q \end{matrix}\middle|\, q\right]_\infty\\
&\times {}_9\phi_8\left[\begin{matrix} b^2/q, b\sqrt{q}, -b\sqrt{q}, bc/q, bd/q, be/q, bf/q, bg/q, bh/q \\ b/\sqrt{q}, -b/\sqrt{q}, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \end{matrix}\middle|\, q;q\right]\\
=\;& \left[\begin{matrix} q, b/q, vq/f, vq/g, vq/h, bf/v, bg/v, bh/v \\ vq, b/v, q^2/f, q^2/g, q^2/h, bf/q, bg/q, bh/q \end{matrix}\middle|\, q\right]_\infty\\
&\times {}_{10}\phi_9\left[\begin{matrix} v, q\sqrt{v}, -q\sqrt{v}, b, vc/q, vd/q, ve/q, f, g, h \\ \sqrt{v}, -\sqrt{v}, vq/b, q^2/c, q^2/d, q^2/e, vq/f, vq/g, vq/h \end{matrix}\middle|\, q;q\right]\\
+\;& \left[\begin{matrix} q,b/q,f,g,h,bq/f,bq/g,bq/h,vc/q,vd/q,ve/q,bq^2/vc,bq^2/vd,bq^2/ve \\ b^2q/v,v/b,q^2/c,q^2/d,q^2/e,q^2/f,q^2/g,q^2/h,bc/q,bd/q,be/q,bf/q,bg/q,bh/q \end{matrix}\middle|\, q\right]_\infty\\
&\times {}_{10}\phi_9\left[\begin{matrix} b^2/v, qb/\sqrt{v}, -qb/\sqrt{v}, b, bc/q, bd/q, be/q, bf/v, bg/v, bh/v \\ b/\sqrt{v}, -b/\sqrt{v}, bq/v, bq^2/vc, bq^2/vd, bq^2/ve, bq/f, bq/g, bq/h \end{matrix}\middle|\, q;q\right],
\end{aligned}
$$

where $v = q^3/cde$ and $q^5 = bcdefgh$. The first term on the left hand side can be manipulated as

$$
\begin{aligned}
&\sum_{k=0}^{\infty}\{1-q^{1+2k}\}\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\middle|\, q\right]_k q^k\\
&= \sum_{k=0}^{\infty}\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\middle|\, q\right]_k q^k\\
&- \sum_{k=0}^{\infty}\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\middle|\, q\right]_k q^{3k+1}\\
&= \sum_{k=0}^{\infty}\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\middle|\, q\right]_k q^k\\
&- \sum_{k=-\infty}^{-1}\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\middle|\, q\right]_{-k-1} q^{-3k-2}\\
&= \sum_{k=0}^{\infty}\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\middle|\, q\right]_k q^k\\
&+ \sum_{k=-\infty}^{-1}\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\middle|\, q\right]_k q^k
\end{aligned}
$$

$$= {}_7\psi_7\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix} \,\middle|\, q; q\right].$$

Hence, we achieve Theorem 4 to finish the proof. $\square$

Taking $b = q^{n+5}/cdefg, h = q^{-n}$ in Theorem 4 and then letting $n \to \infty$, we obtain the following result under the substitutions $e \to b$ and $g \to e$.

Corollary 5. *For* $\vartheta = q^3/bcd$ *and* $\max\{\,|q^4/bcdef|, |q^2/ef|\} < 1$, *there holds the transformation formula between a* ${}_5\psi_5$*-series and a* ${}_8\phi_7$*-series:*

$$\begin{aligned} {}_5\psi_5\left[\begin{matrix} b, c, d, e, f \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f \end{matrix} \,\middle|\, q; \frac{q^4}{bcdef}\right] &= \left[\begin{matrix} q, q^2/ef, \vartheta q/e, \vartheta q/f \\ q^2/e, q^2/f, \vartheta q/ef, \vartheta q \end{matrix} \,\middle|\, q\right]_\infty \\ \times\ {}_8\phi_7\left[\begin{matrix} \vartheta, q\sqrt{\vartheta}, -q\sqrt{\vartheta}, \vartheta b/q, \vartheta c/q, \vartheta d/q, e, f \\ \sqrt{\vartheta}, -\sqrt{\vartheta}, q^2/b, q^2/c, q^2/d, \vartheta q/e, \vartheta q/f \end{matrix} \,\middle|\, q; \frac{q^2}{ef}\right]&. \end{aligned}$$

When $c = q^2/d$, Corollary 5 reduces to (3) under the replacements $e \to c$ and $f \to d$.

Setting $c = q^2/d$ in Theorem 4, we get the following result under the substitutions $g \to c$ and $h \to d$.

Corollary 6. *For* $q^3 = bcdef$, *there holds the transformation formula between a* ${}_5\psi_5$*-series and a* ${}_7\phi_6$*-series:*

$$\begin{aligned} &{}_5\psi_5\left[\begin{matrix} b, c, d, e, f \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f \end{matrix} \,\middle|\, q; q\right] \\ &= \frac{b}{q}\left[\begin{matrix} q, b, c, d, e, f, bq/c, bq/d, bq/e, bq/f \\ b^2, q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, bc/q, bd/q, be/q, bf/q \end{matrix} \,\middle|\, q\right]_\infty \\ &\times\ {}_7\phi_6\left[\begin{matrix} b^2/q, b\sqrt{q}, -b\sqrt{q}, bc/q, bd/q, be/q, bf/q \\ b/\sqrt{q}, -b/\sqrt{q}, bq/c, bq/d, bq/e, bq/f \end{matrix} \,\middle|\, q; q\right] \\ &+ \left[\begin{matrix} q, b/q, q^2/cd, q^2/ce, q^2/cf, q^2/de, q^2/df, q^2/ef \\ q^2/c, q^2/d, q^2/e, q^2/f, bc/q, bd/q, be/q, bf/q \end{matrix} \,\middle|\, q\right]_\infty. \end{aligned}$$

When $e = q^{n+3}/bcd$ and $f = q^{-n}$, Corollary 6 reduces to (5) exactly.

3. Equivalent forms of Theorems 1 and 2

Two ${}_3\psi_3$-series identities, which are respectively due to Bailey [1, Equation 1.2] and Chu and Wang [3, p. 3891], can be expressed as follows:

$${}_3\psi_3\left[\begin{matrix} b, c, d \\ q/b, q/c, q/d \end{matrix} \,\middle|\, q; \frac{q^2}{bcd}\right] = \left[\begin{matrix} q, q/bc, q/bd, q/cd \\ q/b, q/c, q/d, q/bcd \end{matrix} \,\middle|\, q\right]_\infty, \tag{6}$$

$${}_3\psi_3\left[\begin{matrix} b, c, d \\ q^2/b, q^2/c, q^2/d \end{matrix} \,\middle|\, q; \frac{q^4}{bcd}\right] = \frac{-1}{q}\left[\begin{matrix} q, q^2/bc, q^2/bd, q^2/cd \\ q^2/b, q^2/c, q^2/d, q^2/bcd \end{matrix} \,\middle|\, q\right]_\infty, \tag{7}$$

where the convergent conditions for (6) and (7) are $|q/bcd| < 1$ and $|q^2/bcd| < 1$ respectively. The finite forms of them read as

$$\begin{aligned} &{}_5\psi_5\left[\begin{matrix} b, c, d, q^{n+1}/bcd, q^{-n} \\ q/b, q/c, q/d, bcd/q^n, q^{n+1} \end{matrix} \,\middle|\, q; q^2\right] \\ &= \left[\begin{matrix} q, q/bc, q/bd, q/cd \\ q/b, q/c, q/d, q/bcd \end{matrix} \,\middle|\, q\right]_n, \end{aligned} \tag{8}$$

$$
\begin{aligned}
&{}_5\psi_5\left[\begin{matrix} b, c, d, q^{n+3}/bcd, q^{-n} \\ q^2/b, q^2/c, q^2/d, bcd/q^{n+1}, q^{n+2} \end{matrix} \,\middle|\, q; q^3\right] \\
&\quad = \left[\begin{matrix} q^2, q^2/bc, q^2/bd, q^2/cd \\ q^2/b, q^2/c, q^2/d, q^2/bcd \end{matrix} \,\middle|\, q\right]_n \frac{q-1}{q}, \qquad (9)
\end{aligned}
$$

where (8) and (9) are due to Verma and Joshi [9, Equation 3.8] and Chu and Wang [3, Equation 2.8] respectively.

Let k stand for the summation index of the bilateral q-series in (2), (3), (4) and (5). Then there hold the following equivalent relations:

$$
\begin{aligned}
&k \to -k, &&(2) \Rightarrow (6);\\
&k \to -k-1, &&(3) \Rightarrow (7);\\
&k \to -k, &&(4) \Rightarrow (8);\\
&k \to -k-1, &&(5) \Rightarrow (9).
\end{aligned}
$$

Performing the replacement $k \to -k$ in Theorem 1, where k is the summation index of the ${}_7\psi_7$-series, we attain the equivalent form of Theorem 1.

Proposition 7. *For $u = q/cde$ and $q^2 = bcdefgh$, there holds the transformation formula for ${}_7\psi_7$-series:*

$$
\begin{aligned}
&{}_7\psi_7\left[\begin{matrix} b, c, d, e, f, g, h \\ q/b, q/c, q/d, q/e, q/f, q/g, q/h \end{matrix} \,\middle|\, q; q^2\right] \\
&+ \left[\begin{matrix} q, b, c, d, e, f, g, h, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \\ b^2q, 1/b, q/c, q/d, q/e, q/f, q/g, q/h, bc, bd, be, bf, bg, bh \end{matrix} \,\middle|\, q\right]_\infty \\
&\times {}_8\phi_7\left[\begin{matrix} b^2, -qb, bc, bd, be, bf, bg, bh \\ -b, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \end{matrix} \,\middle|\, q; q\right] \\
&= \left[\begin{matrix} q, b, uq/f, uq/g, uq/h, bf/u, bg/u, bh/u \\ uq, b/u, q/f, q/g, q/h, bf, bg, bh \end{matrix} \,\middle|\, q\right]_\infty \\
&\times {}_{10}\phi_9\left[\begin{matrix} u, q\sqrt{u}, -q\sqrt{u}, b, uc, ud, ue, f, g, h \\ \sqrt{u}, -\sqrt{u}, uq/b, q/c, q/d, q/e, uq/f, uq/g, uq/h \end{matrix} \,\middle|\, q; q\right] \\
&+ \left[\begin{matrix} q, b, f, g, h, bq/f, bq/g, bq/h, uc, ud, ue, bq/uc, bq/ud, bq/ue \\ b^2q/u, u/b, q/c, q/d, q/e, q/f, q/g, q/h, bc, bd, be, bf, bg, bh \end{matrix} \,\middle|\, q\right]_\infty \\
&\times {}_{10}\phi_9\left[\begin{matrix} b^2/u, qb/\sqrt{u}, -qb/\sqrt{u}, b, bc, bd, be, bf/u, bg/u, bh/u \\ b/\sqrt{u}, -b/\sqrt{u}, bq/u, bq/uc, bq/ud, bq/ue, bq/f, bq/g, bq/h \end{matrix} \,\middle|\, q; q\right].
\end{aligned}
$$

Taking $b = q^{n+2}/cdefg, h = q^{-n}$ in Proposition 7 and then letting $n \to \infty$, we achieve the following result under the substitutions $e \to b$ and $g \to e$.

Corollary 8. *For $\mu = q/bcd$ and $\max\{|q^2/bcdef|, |q/ef|\} < 1$, there holds the transformation formula between a ${}_5\psi_5$-series and a ${}_8\phi_7$-series:*

$$
\begin{aligned}
&{}_5\psi_5\left[\begin{matrix} b, c, d, e, f \\ q/b, q/c, q/d, q/e, q/f \end{matrix} \,\middle|\, q; \frac{q^3}{bcdef}\right] = \left[\begin{matrix} q, q/ef, \mu q/e, \mu q/f \\ q/e, q/f, \mu q/ef, \mu q \end{matrix} \,\middle|\, q\right]_\infty \\
&\times {}_8\phi_7\left[\begin{matrix} \mu, q\sqrt{\mu}, -q\sqrt{\mu}, \mu b, \mu c, \mu d, e, f \\ \sqrt{\mu}, -\sqrt{\mu}, q/b, q/c, q/d, \mu q/e, \mu q/f \end{matrix} \,\middle|\, q; \frac{q}{ef}\right].
\end{aligned}
$$

When $c = q/d$, Corollary 8 reduces to (6) under the replacements $e \to c$ and $f \to d$.

Setting $c = q/d$ in Proposition 7, we obtain the following result under the substitutions $g \to c$ and $h \to d$.

Corollary 9. *For $q = bcdef$, there holds the transformation formula between a ${}_5\psi_5$-series and a ${}_6\phi_5$-series:*

$$
\begin{aligned}
{}_5\psi_5&\left[\begin{matrix} b, c, d, e, f \\ q/b, q/c, q/d, q/e, q/f \end{matrix}\,\Big|\, q; q^2\right] = {}_6\phi_5\left[\begin{matrix} b^2, -qb, bc, bd, be, bf \\ -b, bq/c, bq/d, bq/e, bq/f \end{matrix}\,\Big|\, q; q\right] \\
&\times\ b\left[\begin{matrix} q, bq, c, d, e, f, bq/c, bq/d, bq/e, bq/f \\ b^2q, q/b, q/c, q/d, q/e, q/f, bc, bd, be, bf \end{matrix}\,\Big|\, q\right]_\infty \\
&+\ \left[\begin{matrix} q, b, q/cd, q/ce, q/cf, q/de, q/df, q/ef \\ q/c, q/d, q/e, q/f, bc, bd, be, bf \end{matrix}\,\Big|\, q\right]_\infty.
\end{aligned}
$$

When $e = q^{n+1}/bcd$ and $f = q^{-n}$, Corollary 9 reduces to (8) exactly.

Employing the replacement $k \to -k-1$ in Theorem 4, where k is the summation index of the ${}_7\psi_7$-series, we get the equivalent form of Theorem 4.

Proposition 10. *For $v = q^3/cde$ and $q^5 = bcdefgh$, there holds the transformation formula for ${}_7\psi_7$-series:*

$$
\begin{aligned}
&{}_7\psi_7\left[\begin{matrix} b, c, d, e, f, g, h \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, q^2/g, q^2/h \end{matrix}\,\Big|\, q; q^3\right] \\
&+\ \frac{b}{q^2}\left[\begin{matrix} q,b,c,d,e,f,g,h,bq/c,bq/d,bq/e,bq/f,bq/g,bq/h \\ b^2,q^2/b,q^2/c,q^2/d,q^2/e,q^2/f,q^2/g,q^2/h,bc/q,bd/q,be/q,bf/q,bg/q,bh/q \end{matrix}\,\Big|\, q\right]_\infty \\
&\times\ {}_9\phi_8\left[\begin{matrix} b^2/q, b\sqrt{q}, -b\sqrt{q}, bc/q, bd/q, be/q, bf/q, bg/q, bh/q \\ b/\sqrt{q}, -b/\sqrt{q}, bq/c, bq/d, bq/e, bq/f, bq/g, bq/h \end{matrix}\,\Big|\, q; q\right] \\
&+\ \frac{1}{q}\left[\begin{matrix} q, b/q, vq/f, vq/g, vq/h, bf/v, bg/v, bh/v \\ vq, b/v, q^2/f, q^2/g, q^2/h, bf/q, bg/q, bh/q \end{matrix}\,\Big|\, q\right]_\infty \\
&\times\ {}_{10}\phi_9\left[\begin{matrix} v, q\sqrt{v}, -q\sqrt{v}, b, vc/q, vd/q, ve/q, f, g, h \\ \sqrt{v}, -\sqrt{v}, vq/b, q^2/c, q^2/d, q^2/e, vq/f, vq/g, vq/h \end{matrix}\,\Big|\, q; q\right] \\
&+\ \frac{1}{q}\left[\begin{matrix} q,b/q,f,g,h,bq/f,bq/g,bq/h,vc/q,vd/q,ve/q,bq^2/vc,bq^2/vd,bq^2/ve \\ b^2q/v,v/b,q^2/c,q^2/d,q^2/e,q^2/f,q^2/g,q^2/h,bc/q,bd/q,be/q,bf/q,bg/q,bh/q \end{matrix}\,\Big|\, q\right]_\infty \\
&\times\ {}_{10}\phi_9\left[\begin{matrix} b^2/v, qb/\sqrt{v}, -qb/\sqrt{v}, b, bc/q, bd/q, be/q, bf/v, bg/v, bh/v \\ b/\sqrt{v}, -b/\sqrt{v}, bq/v, bq^2/vc, bq^2/vd, bq^2/ve, bq/f, bq/g, bq/h \end{matrix}\,\Big|\, q; q\right] = 0.
\end{aligned}
$$

Taking $b = q^{n+5}/cdefg, h = q^{-n}$ in Proposition 10 and then letting $n \to \infty$, we attain the following result under the substitutions $e \to b$ and $g \to e$.

Corollary 11. *For $\vartheta = q^3/bcd$ and $\max\{|q^4/bcdef|, |q^2/ef|\} < 1$, there holds the transformation formula between a ${}_5\psi_5$-series and a ${}_8\phi_7$-series:*

$$
\begin{aligned}
{}_5\psi_5&\left[\begin{matrix} b, c, d, e, f \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f \end{matrix}\,\Big|\, q; \frac{q^6}{bcdef}\right] + \frac{1}{q}\left[\begin{matrix} q, q^2/ef, \vartheta q/e, \vartheta q/f \\ q^2/e, q^2/f, \vartheta q/ef, \vartheta q \end{matrix}\,\Big|\, q\right]_\infty \\
&\times\ {}_8\phi_7\left[\begin{matrix} \vartheta, q\sqrt{\vartheta}, -q\sqrt{\vartheta}, \vartheta b/q, \vartheta c/q, \vartheta d/q, e, f \\ \sqrt{\vartheta}, -\sqrt{\vartheta}, q^2/b, q^2/c, q^2/d, \vartheta q/e, \vartheta q/f \end{matrix}\,\Big|\, q; \frac{q^2}{ef}\right] = 0.
\end{aligned}
$$

When $c = q^2/d$, Corollary 11 reduces to (7) under the replacements $e \to c$ and $f \to d$.

Setting $c = q^2/d$ in Proposition 10, we achieve the following result under the substitutions $g \to c$ and $h \to d$.

Corollary 12. *For $q^3 = bcdef$, there holds the transformation formula between a ${}_5\psi_5$-series and a ${}_7\phi_6$-series:*

$$
\begin{aligned}
&{}_5\psi_5\left[\begin{matrix} b, c, d, e, f \\ q^2/b, q^2/c, q^2/d, q^2/e, q^2/f \end{matrix} \,\Big|\, q; q^3\right] \\
&+ \frac{b}{q^2}\left[\begin{matrix} q, b, c, d, e, f, bq/c, bq/d, bq/e, bq/f \\ b^2, q^2/b, q^2/c, q^2/d, q^2/e, q^2/f, bc/q, bd/q, be/q, bf/q \end{matrix} \,\Big|\, q\right]_\infty \\
&\times {}_7\phi_6\left[\begin{matrix} b^2/q, b\sqrt{q}, -b\sqrt{q}, bc/q, bd/q, be/q, bf/q \\ b/\sqrt{q}, -b/\sqrt{q}, bq/c, bq/d, bq/e, bq/f \end{matrix} \,\Big|\, q; q\right] \\
&+ \frac{1}{q}\left[\begin{matrix} q, b/q, q^2/cd, q^2/ce, q^2/cf, q^2/de, q^2/df, q^2/ef \\ q^2/c, q^2/d, q^2/e, q^2/f, bc/q, bd/q, be/q, bf/q \end{matrix} \,\Big|\, q\right]_\infty = 0.
\end{aligned}
$$

When $e = q^{n+3}/bcd$ and $f = q^{-n}$, Corollary 12 reduces to (9) exactly.

References

[1] W. N. Bailey, On the analogue of Dixon's theorem for bilateral basic hypergeometric series, Quart. J. Math. 1 (1950) 318-320.

[2] W.Y.C. Chen, A.M. Fu, Semi-finite forms of bilateral basic hypergeometric series, Prop. Amer. Math. Soc. 134 (2006) 1719-1725.

[3] W. Chu, C. Wang, Bilateral inversons and terminating basic hypergeometric series identities, Discrete. Math. 309 (2009) 3888-3904.

[4] F. H. Jackson, Certain q-identities, Quart. J. Math. 12 (1941) 167-172.

[5] F. Jouhet, M. Schlosser, Another proof of Bailey's ${}_6\psi_6$ summation, Aequationes Mathematicae 70 (2005) 43-50.

[6] F. Jouhet, Some more Semi-finite forms of bilateral basic hypergeometric series, Ann. Combin. 11 (2007) 47-57.

[7] F. Jouhet, M. Schlosser, New curious bilateral q-series identities, Axioms 1 (2012) 365-371.

[8] G. Gasper, M. Rahman, Basic Hypergeometric Series (2nd edition), Cambridge Univercity Press, Cambridge, 2004.

[9] A. Verma, C. M. Joshi , Some remarks on summation of basic hypergeometric series, Houston J. Math. 5 (1979) 277-294.

[10] C. Zhang, Z. Zhang, Two new transformation formulas of basic hypertgeometric series, J. Math. Anal. Appl. 336 (2007) 777-787.

[11] Z. Zhang, On the bilateral series ${}_5\psi_5$, J. Math. Anal. Appl. 337 (2008) 1002-1009.